

## 2019-2020 IMMUNIZATIONS – DOSES AND INTERVALS

### ENTERING KINDERGARTEN THROUGH GRADE 8

**Note: IL Child Health Exam** dated within 12 months of first day of school is required for Kindergarten and 6<sup>th</sup> Graders.

#### Polio

Kindergarten children immunized according to an all-IPV or all-OPV schedule must show evidence of having received at least 4 doses (as do all children who are given a combination of polio vaccine types), with intervals of no less than 4 weeks apart, and the last dose having been received on/after the 4<sup>th</sup> birthday and at least 6 months since previous dose.

**All other grades:** 3 doses with the last dose qualifying as a booster and received on or after the 4<sup>th</sup> birthday and at least 6 months since previous dose.

#### DTP/DtaP/Td

4 or more doses, with last dose administered on/after the 4<sup>th</sup> birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose.

**For children entering school other than Pre-K, K and 1<sup>st</sup> grade, 3 doses of DTP/DtaP/TD, with last dose administered on/after the 4<sup>th</sup> birthday, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose AND, after the age of 7, one of the doses needs to be a Tdap (Note: others may be TD).**

**Lead Screening & Results** done after 6 months of age.

#### Measles, Mumps and Rubella Vaccine

2 doses, 1<sup>st</sup> dose administered on/after the 1<sup>st</sup> birthday, 2<sup>nd</sup> dose no less than 4 weeks (28 days) after 1<sup>st</sup> dose.

#### \* **Hepatitis B [not required for K-5]**

3 doses needed. The interval of at least 4 weeks (28 days) between 1<sup>st</sup> and 2<sup>nd</sup> dose; the interval of at least 2 months (56 days) between 2<sup>nd</sup> and 3<sup>rd</sup> dose; and the interval between 1<sup>st</sup> and 3<sup>rd</sup> doses of at least 4 months.

#### \*\***Varicella (chickenpox)**

2 doses, with first dose administered on/after the 1<sup>st</sup> birthday and the second dose no less than four weeks (28 days) after the first. Note: The recommended minimum interval between doses is 3 months but can be accepted if at least 4 weeks after the first dose.

#### **Meningococcal Conjugate Vaccine (MCV4) - 6<sup>th</sup>, 7<sup>th</sup> and 8<sup>th</sup> graders:** 1 dose of MCV4 on or after 11 years of age

\* **Hepatitis B** – A 3<sup>rd</sup> dose of hepatitis B vaccine is not required if it can be documented that the child received 2 doses of adult formulation Recombivax-HB vaccine, was 11-15 years of age at the time of vaccine administration, and the interval between receipt of the 2 doses was at least 4 months.

\*\***Varicella** – A physician, health care provider or school health professional can confirm past disease history (in lieu of vaccination) by having examined the infected child, documenting the parent's description of the child's history or reviewing laboratory evidence.

### ENTERING A PROGRAM BELOW THE KINDERGARTEN LEVEL

**Note: IL Child Health Exam** dated within 12 months of enrollment is required. (Must be dated within 6 months for Head Start)

#### Polio

3 or more doses, administered no less than 4 weeks apart.

#### \***DTP/DtaP/Td**

4 or more doses, at intervals of no less than 4 weeks apart, last dose at least 6 months since previous dose.

**Lead Screening & Results** dated anytime after the 1<sup>st</sup> birthday.

**TB Test & Results** dated anytime after the 1<sup>st</sup> birthday.

#### **Measles, Mumps, and Rubella Vaccine**

1 dose, administered on/after the 1<sup>st</sup> birthday.

#### **Hepatitis B**

3 doses needed. The interval of at least 4 weeks (28 days) between 1<sup>st</sup> and 2<sup>nd</sup> dose; the interval of at least 2 months between 2<sup>nd</sup> and 3<sup>rd</sup> dose; and the interval between 1<sup>st</sup> and 3<sup>rd</sup> doses of at least 4 months.

#### **Hib**

Children 24-59 months of age who have not received a primary series according to the Hib schedule, must show proof of receiving 1 dose at 15 months of age or older.

#### \*\***Varicella (chickenpox)**

1 dose, administered on/after the 1<sup>st</sup> birthday.

#### **Pneumococcal Vaccine**

Children 24 – 59 months of age shall show proof of immunization that complies with the pneumococcal conjugate schedule. If they have not received the primary series then they must show proof of receiving 1 dose of the pneumococcal vaccine after 24 months of age.